

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**




Figure 1

Templates for: Dostoevsky, Iyodor

Select a doctor	Select a procedure	<input checked="" type="checkbox"/> Sort by CPT Code
Chaucer, Geoffrey	23130 - ACROMIOPLASTY OR ACROMIONECTOMY, PARTIAL, WITH	49505 - REPAIR INITIAL INGUINAL HERNIA, AGE 5 YEARS OR OVER, REDUCIBLE
Dickens, Charles	49507 - REPAIR INITIAL INGUINAL HERNIA, AGE 5 YEARS OR OVER, REDUCIBLE	
Donne, John	49520 - REPAIR RECURRENT INGUINAL HERNIA, ANY AGE; REDUCIBLE	
Emerson, Ralph Waldo	49521 - REPAIR RECURRENT INGUINAL HERNIA, ANY AGE; INCARCERATED	
Shakespeare, William	49580 - REPAIR UMBILICAL HERNIA, UNDER AGE 5 YEARS; REDUCIBLE	
Tolstoy, Leo	49585 - REPAIR UMBILICAL HERNIA, AGE 5 YEARS OR OVER; REDUCIBLE	
	49587 - REPAIR UMBILICAL HERNIA; AGE 5 YRS OR OVER; REDUCIBLE	
	49650 - LAPAROSCOPY HERNIA REPAIR/INITIAL	
	50398 - CHANGE OF NEPHROSTOMY OR PYELOSTOMY TUBE	

Select a template or templates

Select all Deselect all

Create new template for selected procedure

Edit selected templates

Cancel

Figure 2

Templates for: Dostoevsky, Fyodor

49505 - REPAIR INITIAL INGUINAL HERNIA, AGE 5 YEARS OR OVER; REDUCIBLE

Procedure


Version: Standard template/Inguinal hernia repair

Add [insert text] tags at places in the text where unique case details are to be inserted later.

Select a section	Description of Procedure	View Template
Preoperative Diagnosis	<p>The patient was taken to the operating room. After¹ uneventful administration of general anesthesia via laryngeal mask the patient was placed in the supine position on the operating room table. The patient's abdomen and genitalia was prepped with Betadine solution and draped in the sterile fashion.</p> <p>A transverse skin marking was made approximately one finger breadth above the pubic symphysis and the marked area was infiltrated with 1% Lidocaine solution with 1:100 000 dilution epinephrine.</p> <p>A transverse incision was then made following the skin marking.</p>	
Postoperative Diagnosis		
Anesthesia		
Estimated Blood Loss		
Complications		
Class		
Wound Classification		
Sponge and Needle Counts		
Indications		
Procedure and Findings		
Intraoperative Findings		
Disposition		

Cancel < << >> > Delete... Save

Figure 3



The screenshot shows a software window titled "Create OpNote(s)". The window is divided into several sections:

- Patient Info:** Contains fields for Case # (1234), Surgery Date (11/13/2003), Pt Name (East, Thomas), DOB (12/15/1982), Age (21), and Gender (M).
- Procedure(s):** A text box containing "52005 - CYSTOURETHROSCOPY, WITH URETERAL CATHETERIZATION, WITH OR WITHOUT IRRIGATION, INSTILLATION, OR URETEROPYELOGRAPHY, EXCLUSIVE OF RADIOLOGIC SERVICE".
- Surgeon(s):** A text box containing "Fyodor Dostoevsky, MD".
- Left Sidebar:** Contains two buttons: "Apply template" (with a document icon) and "Cycle through [insert text] tags" (with a circular arrow icon).
- Select a section:** A list box containing the following items: "Preoperative Diagnosis", "Postoperative Diagnosis", "Anesthesia", "Estimated Blood Loss", "Complications", "Class", "Wound Classification", "Sponge and Needle Counts", "Indications", "Description of Procedure", "Intraoperative Findings", and "Disposition". The "Preoperative Diagnosis" item is currently selected.
- Preoperative Diagnosis:** A large, empty text box for entering the preoperative diagnosis.
- Bottom:** Contains a "Cancel" button on the left, a set of four small navigation buttons in the center, and "Print ..." and "Save" buttons on the right.

Figure 5

Apply Template

Procedures

☐ Sort by CPT Code

52214 - CYSTOURETHROSCOPY, WITH FULGURATION (INCLUDING CRYOSU
52224 - CYSTOURETHROSCOPY, WITH FULGURATION (INCLUDING CRYOSU
52332 - CYSTOURETHROSCOPY, WITH INSERTION OF INDWELLING URETEF
52282 - CYSTOURETHROSCOPY, WITH INSERTION OF URETHRAL STENT
52270 - CYSTOURETHROSCOPY, WITH INTERNAL URETHROTOMY; FEMALE
52275 - CYSTOURETHROSCOPY, WITH INTERNAL URETHROTOMY; MALE
52315 - CYSTOURETHROSCOPY, WITH REMOVAL OF FOREIGN BODY, CALCI
52310 - CYSTOURETHROSCOPY, WITH REMOVAL OF FOREIGN BODY, CALCI
52277 - CYSTOURETHROSCOPY, WITH RESECTION OF EXTERNAL SPHINCTE
52005 - CYSTOURETHROSCOPY, WITH URETERAL CATHETERIZATION, WITH
OR WITHOUT IRRIGATION, INSTILLATION, OR URETEROPYELOGRAPHY,
EXCLUSIVE OF RADIOLOGIC SERVICE;

Templates

Standard Template/MALE Cystoscopy w/bilateral retrograde pyelograms / Las
Standard Template/FEMALE Cystoscopy w/bilateral retrograde pyelograms / Las

Cancel

Apply

Figure 6

View and Print OpNote

MainReport

Operative Report

Case Number: 1,593 Patient Acc# Number: 1,191 Chart Number: N/A
Patient Name: East, Thomas DOB: 12/15/82 Age: 21 Sex: M

Surgery Date: 11/13/03

Text Object: Yodanis Dostoevsky, MD

Procedure(s): 52005 - CYSTOURETHROSCOPY, WITH URETERAL CATHETERIZATION, WITH OR WITHOUT IRRIGATION, INSTILLATION, OR URETEROPYELOGRAPHY, EXCLUSIVE OF RADIOLOGIC SERVICE;

PREOPERATIVE DIAGNOSIS: Microscopic hematuria.

POSTOPERATIVE DIAGNOSIS: Same.

ANESTHESIA: General.

COMPLICATIONS: None.

INDICATIONS: This is a 21 year old white male who was found to have something or other. He was scheduled for cystoscopy and bilateral retrograde pyelogram to complete the evaluation for a problem of some kind.

DESCRIPTION OF PROCEDURE: The patient was taken to the operating room. After uneventful administration of general anesthesia, he was placed in a lithotomy position on the operating table. The genitalia was prepped with Betadine solution and draped in a sterile fashion. A 22 French semi-rigid cystoscope was introduced. The urethra was normal. The prostate gland was normal and without enlargement. The bladder was entered and carefully inspected. Both ureteral orifices were visualized and palpated. The bladder mucosa was normal. The bladder wall was [insert text]. Bilateral retrograde pyelogram was next performed using a 58 cm-tip ureteral catheter. Approximately 5-6 cc of contrast material was first injected into the [insert text] upper collecting system. Fluoroscopy images were obtained. The [insert text] upper collecting system was next visualized and contrast injected. The [insert text] upper collecting system was next visualized and contrast injected. The [insert text] upper collecting system was next visualized and contrast injected.

Current Page No: 1 Total Page No: 1 Zoom Factor: Page Width

Figure 7




Figure 8